

Developmental rates of bovine nuclear transfer embryos derived from different fetal non transfected and transfected cells

Ricardo Felmer D.¹ ✉ · María Elena Arias¹

¹ Laboratorio de Biotecnología Animal, Instituto de Investigaciones Agropecuarias, INIA - Carillanca, Temuco, Chile

✉ Corresponding author: rfelmer@inia.cl
Received December 12, 2010 / Accepted March 14, 2011
Published online: May 15, 2011
© 2011 by Pontificia Universidad Católica de Valparaíso, Chile

Abstract Since the first successful somatic cell nuclear transfer (SCNT) experiments were carried out, a number of domestic and agriculture species have been cloned using donor cells derived from different sources and origin. However, differences in nuclear transfer efficiency both *in vitro* and *in vivo* have been generally observed. These differences may be accentuated when transgenic cell lines are used as nuclear donors in an attempt to generate transgenic cloned offspring. The present study examined the suitability of cell lines derived from 3 different fetal sources and the effects of genetic manipulation of donor fetal fibroblasts with a red fluorescent plasmid, on the *in vitro* developmental potential and quality of nuclear transfer derived bovine embryos. We observed no differences in the cleavage rate of nuclear transfer embryos generated with any of the cell lines evaluated. However, the blastocyst rate was significantly affected when cell lines were derived from the 3 different fetal sources (21, 18 and 11%, respectively) or from 2 transgenic clonal cell lines that had originated from the same primary fetal cell (18 and 10%, respectively). Despite this difference, quality of embryos as measured by the total number of cells and by assessing some morphology aspects of their appearance was not different. Together these results indicate that fetal fibroblast cell lines derived from different fetal sources and transgenic clonal cell lines that had originated from the same fetus results in different *in vitro* developmental potential when used as donors for nuclear transfer experiments. Further studies, including evaluation of pregnancy rates, development to term, and epigenetic modifications of these cell lines will be necessary to better understand the differences observed in nuclear transfer efficiency.

Keywords: bovine embryos, fetal fibroblasts, nuclear transfer, transfected cells

INTRODUCTION

Cloning by somatic cell nuclear transfer (SCNT) has been successfully achieved in a number of domestic and agriculture species (for a comprehensive review see Campbell et al. 2007). A range of embryonic, fetal or adult cells have been used as nuclear donors, including mammary gland cells, cumulus cells, oviductal cells, leukocytes, granulosa cells, germ cells, liver cells and muscle cells (Brem and Kühholzer, 2002). However, it is still unclear which cell type is the most successful for nuclear transfer into oocytes and despite the success with these cells, different efficiencies of *in vitro* and *in vivo* development of nuclear transfer embryos have been generally observed (Zakhartchenko et al. 1999; Inoue et al. 2003; Poehland et al. 2007). For instance, when the efficiencies of various cell types from adult, newborn and fetal donor cells were compared in cattle, no significant difference were found in the percentage of embryos reaching to the blastocyst stage (Kato et al. 2000). Similar results were also observed in mice when macrophages, fibroblasts and spleen cells were used (Wakayama and Yanagimachi, 2001). However, a number of studies have also shown significant differences depending on the cell type source, age and/or sex of the donor cell used for nuclear transfer (Lagutina et al. 2005). Nuclear transfer technology also allows the opportunity for animal genetic manipulation (Clark, 1998; Houdebine, 2009). In order to do this, nuclear donor cells must be genetically manipulated *in vitro* to target the desired modification in the animal. Therefore, an extended period of culture is required for selection of transgenic cells resistant to the marker gene. This extended period of

culture may have profound effects on the developmental potential of transgenic cells used as nuclear donors as described by a number of reports (Zakhartchenko et al. 2001; Arat et al. 2002; Iguma et al. 2005; Wang et al. 2007). Therefore, in the present study we examined the effect of cell lines derived from different fetal sources and genetic manipulation of these cells, on the *in vitro* developmental potential and quality of nuclear transfer derived bovine embryos in order to establish if donor source, transfection and longer culture of these cells, could have an effect on the developmental potential of nuclear transfer embryos.

MATERIALS AND METHODS

Derivation and transfection of donor cells

Bovine fetal fibroblasts (BFF) were isolated from 3 different Holstein female 50-70 days old fetuses, recovered at a local slaughterhouse. Cell explants were cultured in Dulbecco's Modified Eagle's Medium (DMEM, Gibco) supplemented with 10% FCS (HyClone) at 38.5°C in a humidified atmosphere of 5% CO₂. When cells reached confluency, they were collected by trypsinization and frozen for future nuclear transfer experiments. A primary cell line derived from the female fetus 1 (BFF 1) was used for transfection experiments on passage 5. Two µg of a linearized plasmid (*Afl* II digested) containing the red fluorescence protein marker (pHcRed1-nuc; Clontech) was combined with Lipofectamine (Invitrogen), according to the manufacturer instructions. Cells were selected with 800 µg/ml Geneticin (G418, HyClone) for 7-10 days and single colonies isolated by ring cloning. A portion of cells was used for DNA isolation, confirmation of construct integration by PCR and chromosome analysis. Remaining cells were frozen for future nuclear transfer experiments.

Oocyte collection, maturation and nuclear transfer

Bovine ovaries were collected at a local slaughterhouse (Frigorifico Temuco) and grade 1 cumulus oocyte complexes (COCs) were selected for maturation, that was carried out in Medium 199 (Sigma) supplemented with 10% FCS (HyClone) and hormones at 38.5°C in a humidified atmosphere of 5% CO₂ in air. After 17 hrs of maturation, oocytes were stained with Hoechst 33342 (Sigma) and enucleated by aspiration of the MII plate using an inverted microscope (Nikon TS100F) and Narishige micromanipulators. Prior to nuclear transfer, donor cells (transfected and non-transfected) were grown to confluency for 5 days to induce quiescence. These cells were microsurgically placed into the perivitelline space evacuated during enucleation and cell-cytoplasm complexes were fused in sorbitol media with a single DC pulse of 170 volts/mm and 15 µseconds delivered by an Electrocell Manipulator 830 (BTX). Activation was carried out with 5 µM Ionomycin for 5 min followed by incubation in KSOM (Millipore) media containing cycloheximide (10 µg/ml, Sigma) and cytochalasin B (5 µg/ml, Sigma) for 5 hrs. After activation, the nuclear transfer units were cultured in 30 µl drops of KSOM media under mineral oil at 38.5°C and at 5% CO₂, 5% O₂, 90% N₂. On day 3 (nuclear transfer = day 0) cleavage rate was recorded and embryos were transferred to KSOM+5% FCS and cultured until day 7, time when blastocyst rate and total number of cells were recorded. All the cell lines were evaluated on the same day and the experiments were replicated 4 times.

Cell number count and statistical analysis

Cell number of blastomeres in day 7 blastocysts was scored by incubating embryos in media (TCM-199) containing 10 µg/ml bisbenzamide in absolute ethanol at room temperature for 10 min. Blastocysts were then treated in 50% (v/v) glycerol, mounted onto a glass slide, and visualized under epifluorescent microscope coupled with a Filter UV-2E/C DAPI. Data analysis for embryo development and cell count was carried out by descriptive statistics based on average and standard error calculated for each of the variables, using Stat graphics Plus 5.1 Software. One-way ANOVA was used to test for statistically significant differences ($P < 0.05$) among groups for cleavage, blastocysts rate and cell counting. In cases where statistically significant differences were observed, the least significant difference test (LSD) was invoked to determine where those differences existed.

RESULTS

Bovine fetal fibroblast (BFF) cell lines were successfully established from all 3 fetuses recovered at the slaughterhouse (described herein as BFF1, BFF2 and BFF3, respectively). Cell lines were cultured up to passage 4 and nuclear transfer experiments were carried out with these cells on passage 5-6. Stably transfected cells were also generated with the marker gene pHcRed1-nuc, allowing selection both by resistance to neomycin and the intensity of red fluorescence protein signal (Figure 1). Fifteen neomycin resistant cell lines were generated, 5 of which expressed red fluorescent protein at a higher level. Karyotyping analysis of these cells showed a normal chromosome complement ($2n = 60$; XX) in 4 out of 5 cell lines analyzed. This analysis allowed the selection of 2 normal transfected cell lines for further nuclear transfer experiments (Figure 1; clones BFF-Tg 1.1 and BFF-Tg 1.2, respectively), that were used on passage 9-10. The assessment of transgenic cloned embryos expressing red fluorescent protein during the embryonic development up to the blastocyst stage, was described previously (Felmer and Arias, 2008). The efficiency of nuclear transfer generated bovine embryos was compared using these transfected and non-transfected bovine fetal fibroblast cell lines on independent replicates. No significant difference in terms of cleavage rate was observed between embryos generated for any of the transfected and non-transfected cell lines (59 to 64%). However, a significant difference in the rate of blastocyst generated from cell lines derived from the non-transfected and/or transfected group was observed (Table 1). One transfected (BFF tg 1.2) and one non-transfected (BFF 3) cell line had the lowest rate of blastocyst (10 and 11%, respectively), while transfected (BFF tg 1.1) and non-transfected (BFF 1 and 2) cell lines had the best developmental capability (21, 18 and 18%, respectively). The quality of nuclear transfer embryos generated by these cells, as measured by the total number of cells and by assessing some morphology aspects of their appearance, was not different for any of the cell lines used as nuclear donors.




Fig. 1 Transgenic bovine fetal fibroblast (BFF tg 1.1 and 1.2) in culture. (a) and (d): Confluent transfected bovine fetal fibroblasts derived from fetus N°1 (cell lines BFF tg 1.1 and BFF tg 1.2, respectively). (b) and (e): Epifluorescent image of transfected cell lines BFF tg 1.1 and BFF tg 1.2, respectively. (c) and (f): Representative karyogram of transfected cell lines BFF tg 1.1 and BFF tg 1.2, respectively. Magnification 100 X.

DISCUSSION

In this report, we investigated the suitability of cell lines derived from 3 different fetal sources and the effects of genetic manipulation of these cells on the *in vitro* developmental potential of nuclear transfer derived bovine embryos. We chose to use fetal cells as donors for nuclear transfer experiments, because these cells are believed to have less genetic damage and better proliferative ability than adult

somatic cells and therefore are more robust for transfection experiments. We observed a significant effect between cell lines derived from different fetal sources despite the fact that all cell lines belonged to the same breed of origin, were isolated within the same day and were cultivated under equal culture conditions. Furthermore, all the cell lines were at the same passage number by the time of nuclear transfer experiments were carried out (between 5-6 passages) and were subjected to the same cell cycle synchronization protocol, as described in Material and Methods section. Although there was no difference in the cleavage rate for any of the fetal fibroblast cell lines used, a significant difference was observed in the rate of blastocyst that was consistently higher for 2 of the 3 fetal fibroblast cell lines assessed (21, 18 and 11% for BFF 1, BFF 2 and BFF 3, respectively).

Table 1. *In vitro* development of bovine NT embryos derived from 3 different fetal sources and from 2 transfected cell lines derived from 1 fetus.

Cell Line	Nº of embryos cultured	Cleavage <i>n</i> (%)	Blastocyst <i>n</i> (%)	number of cells (mean ± SD)
BFF 1	118	76 (64)	25 (21) ^a	95 ± 9.1
BFF 2	119	72 (61)	21 (18) ^a	81 ± 11.1
BFF 3	119	70 (59)	13 (11) ^b	84 ± 5.8
BFF tg 1.1	115	73 (64)	20 (18) ^a	85 ± 12.0
BFF tg 1.2	119	72 (61)	12 (10) ^b	83 ± 14.5

BFF: Non-transfected bovine fetal fibroblast cell lines derived from different fetuses. BFF tg: Transfected bovine fetal fibroblast clonal cell lines derived from fetus 1. Experiments represent 4 replicates. Cleavage, blastocyst and nuclei stain were registered at 72, 168 and 168 hrs, respectively. Data followed by different letters are significantly different ($P < 0.05$).

Results described previously using different primary cell lines of adult bovine somatic cells indicated that the primary donor cell culture affects *in vitro* blastocyst development, initial pregnancy rates, and the percentage of live births (Miyoshi et al. 2003; Poehland et al. 2007). Similar results were obtained in pig cloning studies, where not only different cell lines, but also different clones derived from one primary cell line used as nuclear donors resulted in different development beyond day 90 of gestation (Kühholzer et al. 2001) and in horse cloning studies where cumulus, granulose, fetal and adult fibroblast cells had all *in vitro* and *in vivo* differential developmental potential (Lagutina et al. 2005). However, contradictory results have also been observed when the efficiency of nuclear transfer using a number of somatic cell types from adult, newborn, and fetal female and male donor cattle were compared with no difference in the percentage of embryos that developed to the blastocyst stage being reported (Kato et al. 2000). Similar results were also observed using various cell types derived from mice of different strains, sexes and ages (Wakayama and Yanagimachi, 2001). This marked difference observed for any of the cell types and/or cell lines used in nuclear transfer experiments may be due to epigenetic effects, because even within a primary cell culture, the generation of cell lines from that culture showed some lines more suitable than others for nuclear transfer (Kühholzer et al. 2001). Modifications that may occur during cell culture might result in epigenetic changes to the genomes making these cells more or less capable of being reprogrammed. In fact, wide-spread abnormalities in gene expression in embryos and cloned animals are beginning to emerge, which suggests that transcriptional regulation mechanisms were impaired possibly at different levels during the reprogramming process (Dean et al. 2001; Beyhan et al. 2007; Suzuki et al. 2008).

Transgenic cloned offspring has been produced by SCNT in sheep, cows and other species (Schnieke et al. 1997; Cibelli et al. 1998; Bordignon et al. 2003; Echelard et al. 2009). The generation of transgenic animals by this approach requires the genetic manipulation of donor cells, a process that has been observed to cause a dramatic effect on the efficiency of the nuclear transfer technique and in some cases in the failure of generating live offspring (Iguma et al. 2005; Jang et al. 2006). This situation rise the question of whether the efficiency of SCNT in generating transgenic animals would be affected by the transgene itself or the modified donor cells. In this report, we carried out cytogenetic analysis of transfected BFF cells expressing a red fluorescent protein, since it has been reported a direct relationship between chromosomal anomalies in nuclear transfer embryos and in their respective nuclear donor cells (Bureau et al. 2003). A clonal cell line with abnormal chromosome complement was found within 5 clonal cell lines analyzed, which allowed to exclude this cell line for future nuclear transfer analysis. Remaining 4 clonal cell lines, had all normal chromosome complement ($2n = 60; XX$). Two of these transfected clonal cell lines were chosen for nuclear transfer evaluation. These cell lines (BFF tg 1.1 and BFF tg 1.2, respectively), which derived from the same primary culture (BFF 1), had a significantly different developmental rate to the blastocyst stage (18 v/s 10%, respectively), although

cleavage rate was not affected (64 and 61%, respectively). Interestingly, one of the transfected cell lines (BFF tg 1.1) showed no differences in the cleavage (64 and 64%, respectively) and/or blastocyst rate (18 and 21%, respectively), when compared to the primary culture from which it was derived (BFF 1), while the other one (BFF tg 1.2) had a significant decrease in the rate of blastocyst (10 v/s 21%, respectively). This result indicates that transfection and longer passage number of transfected cells not necessarily affect the nuclear transfer efficiency *in vitro*, but this effect is more dependent on each clonal cell line, due to differences in the site of integration, number of integrated transgene copies, and/or other epigenetic effects as result of the *in vitro* manipulation. It was previously observed a significant difference in the efficiency in generating nuclear transfer embryos when transfected ear fibroblasts were used as compared to the non-transfected counterpart, regardless of passage number (Bhuiyan et al. 2004). However, contradictory results were observed by others, with no differences in fusion, cleavage, blastocyst or pregnancy rates between transfected and non-transfected ear fibroblasts (Iguma et al. 2005). This discrepancy with transfected cells, has been suggested to be due to differences in vector type and transfection protocols (Bhuiyan et al. 2004), donor cell culture conditions (Wells et al. 2003), and the transgene and/or site of transgene integration (Hodges and Stice, 2003). In fact, a recent study with transfected cells harboring different foreign genes (*hFIX*, *hALB* and *hTF*), confirmed this observation since significant differences in fusion, cleavage and/or blastocyst rate was observed when these cells were used in nuclear transfer experiments (Fu et al. 2008).

In conclusion, the results presented here suggest that cloning efficiency is not only affected by the source of cells used as nuclear donors. Variations in nuclear transfer results between experiments and between different cell lines as described here and elsewhere, are probably due to deviations in the epigenetic regulation of the gene expression profile of each donor cell line, as result of genetic disturbances resulting from the culture and/or manipulation conditions. Therefore, whether transfected and non-transfected cells are used, it is necessary first to assess the *in vitro* developmental potential of each donor cell line in order to select those with the highest rate of blastocyst development and thus the most suitable for nuclear transfer experiments, bearing in mind that a high rate of blastocyst *in vitro* will not necessarily correlate with implantation and normal development of these embryos after transfer. Further studies, including evaluation of pregnancy rates, development to term, and epigenetic modifications of these cell lines will be necessary to better understand the differences observed in nuclear transfer efficiency.

Financial support: This research was supported by grants from FONDECYT 1080216 and FONDEF D0311074

REFERENCES

- ARAT, S.; GIBBONS, J.; RZUCIDLO, S.J.; RESPESS D.S.; TUMLIN, M. and STICE, S.L. (2002). *In vitro* development of bovine nuclear transfer embryos from transgenic clonal lines of adult and fetal fibroblast cells of the same genotype. *Biology of Reproduction*, vol. 66, no. 6, p. 1768-1774. [\[CrossRef\]](#)
- BEYHAN, Z.; FORSBERG, E.J.; EILERTSEN, K.J.; KENT-FIRST, M. and FIRST, N.L. (2007). Gene expression in bovine nuclear transfer embryos in relation to donor cell efficiency in producing live offspring. *Molecular Reproduction and Development*, vol. 74, no. 1, p. 18-27. [\[CrossRef\]](#)
- BHUIYAN, M.M.U.; CHO, J.; JANG, G.; PARK, E.; KANG, S.; LEE, B. and HWANG, W. (2004). Effect of transfection and passage number of ear fibroblasts on *in vitro* development of bovine transgenic nuclear transfer embryos. *Journal of Veterinary Medical Science*, vol. 66, no. 3, p. 257-261. [\[CrossRef\]](#)
- BORDIGNON, V.; KEYSTON, R.; LAZARIS, A.; BILODEAU, A.S.; PONTES, J.H.; ARNOLD, D.; FECTEAU, G.; KEEFER, C. and SMITH, L.C. (2003). Transgene expression of green fluorescent protein and germ line transmission in cloned calves derived from *in vitro*-transfected somatic cells. *Biology of Reproduction*, vol. 68, no. 6, p. 2013-2023. [\[CrossRef\]](#)
- BREM, G. and KÜHHOLZER, B. (2002). The recent history of somatic cloning in mammals. *Cloning and Stem Cells*, vol. 4, no. 1, p. 57-63. [\[CrossRef\]](#)
- BUREAU, W.S.; BORDIGNON, V.; LEVEILLEE, C.; SMITH, L.C. and KING, W.A. (2003). Assessment of chromosomal abnormalities in bovine nuclear transfer embryos and in their donor cells. *Cloning and Stem Cells*, vol. 5, no. 2, p. 123-132. [\[CrossRef\]](#)
- CAMPBELL, K.H.S.; FISHER, P.; CHEN, W.C.; CHOI, I.; KELLY, R.D.W.; LEE J.T.H. and XHU, J. (2007). Somatic cell nuclear transfer: Past, present and future perspectives. *Theriogenology*, vol. 68, supp. 1, p. S214-S231. [\[CrossRef\]](#)
- CIBELLI, J.B.; STICE, S.L.; GOLUEKE, P.J.; KANE, J.J.; JERRY, J.; BLACKWELL, C.; PONCE DE LEON, F.A. and ROBL, J.M. (1998). Cloned transgenic calves produced from nonquiescent fetal fibroblasts. *Science*, vol. 280, no. 5367, p. 1256-1258. [\[CrossRef\]](#)
- CLARK, A.J. (1998). Gene expression in the mammary glands of transgenic animals. *Biochemical Society Symposia*, vol. 63, p. 133-140.

- DEAN, W.; SANTOS, F.; STOJKOVIC, M.; ZAKHARTCHENKO, V.; WALTER, J.; WOLF, E. and REIK, W. (2001). Conservation of methylation reprogramming in mammalian development: Aberrant reprogramming in cloned embryos. *Proceedings of the National Academy of Sciences of the United States of America*, vol. 98, no. 24, p. 13734-13738. [\[CrossRef\]](#)
- ECHELARD, Y.; WILLIAMS, J.L.; DESTREMPES, M.M.; KOSTER, J.A.; OVERTON, S.A.; POLLOCK, D.P.; RAPIEJKO, K.T.; BEHBOODI, E.; MASIELLO, N.C.; GAVIN, W.G.; POMMER, J.; VAN PATTEN, S.M.; FABER, D.C.; CIBELLI, J.B. and MEADE, H.M. (2009). Production of recombinant albumin by a herd of cloned transgenic cattle. *Transgenic Research*, vol. 18, no. 3, p. 361-376. [\[CrossRef\]](#)
- FELMER, R. and ARIAS, M.E. (2008). Advances in the development of a noninvasive embryo model for the evaluation of the quality of cloned embryos subjected to different treatments. *Electronic Journal of Biotechnology*, vol. 11, no. 5. [\[CrossRef\]](#)
- FU, J.; GUAN, P.; ZHAO, L.; LI, H.; HUANG, S.; ZENG, F. and ZENG, Y. (2008). Effects of donor cells on *in vitro* development of cloned bovine embryos. *Journal of Genetics and Genomics*, vol. 35, no. 5, p. 273-278. [\[CrossRef\]](#)
- HODGES, C.A. and STICE, S.L. (2003). Generation of bovine transgenics using somatic cell nuclear transfer. *Reproductive Biology and Endocrinology*, vol. 1, no. 1-7, p. 1-81. [\[CrossRef\]](#)
- HOUEBINE, L.M. (2009). Production of pharmaceutical proteins by transgenic animals. *Comparative Immunology, Microbiology and Infectious Diseases*, vol. 32, no. 2, p. 107-121. [\[CrossRef\]](#)
- IGUMA, L.T.; LISKAUSKAS, S.F.; MELO, E.O.; FRANCO, M.M.; PIVATO, I.; VIANNA, G.R.; SOUSA, R.V.; DODE, M.A.; ARAGÃO, F.J.; RECH, E.L. and RUMPF, R. (2005). Development of bovine embryos reconstructed by nuclear transfer of transfected and non-transfected adult fibroblast cells. *Genetics and Molecular Research*, vol. 4, no. 1, p. 55-66.
- INOUE, K.; OGONUKI, N.; MOCHIDA, K.; YAMAMOTO, Y.; TAKANO, K.; KOHDA, T.; ISHINO, F. and OGURA, A. (2003). Effects of donor cell type and genotype on the efficiency of mouse somatic cell cloning. *Biology of Reproduction*, vol. 69, no. 4, p. 1394-1400. [\[CrossRef\]](#)
- JANG, G.; BHUIYAN, M.M.U.; JEON, H.Y.; KO, K.H.; PARK, H.J.; KIM, M.K.; KIM, J.J.; KANG, S.K.; LEE, B.C. and HWANG, W.S. (2006). An approach for producing transgenic cloned cows by nuclear transfer of cells transfected with human alpha 1-antitrypsin gene. *Theriogenology*, vol. 65, no. 9, p. 1800-1812. [\[CrossRef\]](#)
- KATO, Y.; TANI, T. and TSUNODA, Y. (2000). Cloning of calves from various somatic cell types of male and female adult, newborn and fetal cows. *Journal of Reproduction and Fertility*, vol. 120, no. 2, p. 231-237. [\[CrossRef\]](#)
- KÜHHOLZER, B.; HAWLEY, R.J.; LAI, L.; KOLBER-SIMONDS, D. and PRATHER, R.S. (2001). Clonal lines of transgenic fibroblast cells derived from the same fetus result in different development when used for nuclear transfer in pigs. *Biology of Reproduction*, vol. 64, no. 6, p. 1695-1698. [\[CrossRef\]](#)
- LAGUTINA, I.; LAZZARI, G.; DUCHI, R.; COLLEONI, S.; PONDERATO, N.; TURINI, P.; CROTTI, G. and GALLI, C. (2005). Somatic cell nuclear transfer in horses: Effect of oocyte morphology, embryo reconstruction method and donor cell type. *Reproduction*, vol. 130, no. 4, p. 559-567. [\[CrossRef\]](#)
- MIYOSHI, K.; RZUCIDLO, S.J.; PRATT, S.L. and STICE, S.L. (2003). Improvements in cloning efficiencies may be possible by increasing uniformity in recipient oocytes and donor cells. *Biology of Reproduction*, vol. 68, no. 4, p. 1079-1086. [\[CrossRef\]](#)
- POEHLAND, R.; AL-ROSTUM, F.; BECKER, F.; VIERGUTZ, T.; BRUNNER, R.M.; KANITZ, W. and BHOJWANI, S. (2007). Donor cell lines considerably affect the outcome of somatic nuclear transfer in the case of bovines. *Journal of Reproduction and Development*, vol. 53, no. 4, p. 737-748. [\[CrossRef\]](#)
- SCHNIEKE, A.E.; KIND, A.J.; RITCHIE, W.A.; MYCOCK, K.; SCOTT, A.R.; RITCHIE, M.; WILMUT, I.; COLMAN, A. and CAMPBELL, K.H. (1997). Human factor IX transgenic sheep produced by transfer of nuclei from transfected fetal fibroblasts. *Science*, vol. 278, no. 5346, p. 2130-2133. [\[CrossRef\]](#)
- SUZUKI, T.; KONDO, S.; WAKAYAMA, T.; CIZDZIEL, P.E. and HAYASHIZAKI, Y. (2008). Genome-wide analysis of abnormal H₃K₉ acetylation in cloned mice. *PLoS One*, vol. 3, no. 4, p. e1905. [\[CrossRef\]](#)
- WAKAYAMA, T. and YANAGIMACHI, R. (2001). Mouse cloning with nucleus donor cells of different age and type. *Molecular Reproduction and Development*, vol. 58, no. 4, p. 376-383. [\[CrossRef\]](#)
- WANG, H.; AO, H.; PAN, Q.; LI, R.; ZHAO, M.; LIAN, Z.; LI, N. and WU, C. (2007). Effects of different states of sheep fetal fibroblasts as donor cells on the early development *in vitro* of reconstructed sheep embryos. *Science in China Series C: Life Sciences*, vol. 50, no. 2, p. 178-185. [\[CrossRef\]](#)
- WELLS, D.N.; LAIBLE, G.; TUCKER, F.C.; MILLER, A.L.; OLIVER, J.E.; XIANG, T.; FORSYTH, J.T.; BERG, M.C.; COCKREM, K.; L'HUILLIER, P.J.; TERVIT, H.R. and OBACK, B. (2003). Coordination between donor cell type and cell cycle stage improves nuclear cloning efficiency in cattle. *Theriogenology*, vol. 59, no. 1, p. 45-59. [\[CrossRef\]](#)
- ZAKHARTCHENKO, V.; ALBERIO, R.; STOJKOVIC, M.; PRELLE, K.; SCHERNTHANER, W.; STOJKOVIC, P.; WENIGERKIND, H.; WANKE, R.; DÜCHLER, M.; STEINBORN, R.; MUELLER, M.; BREM, G. and WOLF, E. (1999). Adult cloning in cattle: Potential of nuclei from a permanent cell line and from primary cultures. *Molecular Reproduction and Development*, vol. 54, no. 3, p. 264-272. [\[CrossRef\]](#)
- ZAKHARTCHENKO, V.; MUELLER, S.; ALBERIO, R.; SCHERNTHANER, W.; STOJKOVIC, M.; WENIGERKIND, H.; WANKE, R.; LASSNIG, C.; MUELLER, M.; WOLF, E. and BREM, G. (2001). Nuclear transfer in cattle with non-transfected and transfected fetal or cloned transgenic fetal and postnatal fibroblasts. *Molecular Reproduction and Development*, vol. 60, no. 3, p. 362-369. [\[CrossRef\]](#)

How to reference this article:

FELMER, R. and ARIAS, M.E. (2011). Developmental rates of bovine nuclear transfer embryos derived from different fetal non transfected and transfected cells. *Electronic Journal of Biotechnology*, vol. 14, no. 3. <http://dx.doi.org/10.2225/vol14-issue3-fulltext-8>